

Aim to perform this programme a minimum of once per day unless prescribed otherwise. As with any new exercise, start slowly (repetitions as able) and build up as you are able within the guidelines below. Pain should not exceed 5/10 whilst completing this exercise programme.




1. Single leg pendulum

- Standing with feet shoulder-width apart, take your weight onto one leg, bringing your other knee up level with the hip.
- From this position, keep your weight on the standing leg, push your other foot down and out backwards like you are pressing a wall away until it is straight out behind you whilst simultaneously bending your torso forward to counterbalance yourself.
- Try to return to the start without putting your foot down to the floor. It is okay if you cannot do this, but practice will improve it. Repeat on both legs.
- **Perform this exercise up to 15 times, rest for 1 minute.**
- **Repeat routine 2 to 3 times.**



2. Step and reach

- Start by standing with all the weight on one leg.
- Rest the other foot next to the standing foot for balance.
- Now with the non-standing leg, reach out sideways (hovering just above the floor) as far as possible whilst at the same time bending the standing leg.
- Then return to the start position, maintaining your balance.
- Repeat in a forwards and backwards direction, and continue to do this in a cycle of the three movements, only touching down if you start to lose your balance.
- **Perform this exercise up to 15 times, rest for 1 minute.**
- **Repeat routine 2 to 3 times.**



3. Bridge with leg lift

- Lie on your back on the floor or bed with your knees bent at 90 degrees and your feet on the floor and your arms down by your side.
- Push your heels down into the floor to gently lift your hips upwards. Lift as high as you feel comfortable but no higher than a straight line between hips, knees, and shoulders.
- If you feel discomfort, then try squeezing your tummy and bum muscles before starting the lift.
- From this position, lift one leg slightly off the floor and return.
- Repeat on the opposite side.
- Try and keep the pelvis still.
- **Perform this exercise up to 15 times, rest for 1 minute.**
- **Repeat routine 2 to 3 times.**